

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/239846874>

# First record of Fraser's dolphin *Lagenodelphis hosei* for the Dutch Caribbean

Article in *Marine Biodiversity Records* · May 2012

DOI: 10.1017/S1755267212000279

CITATIONS

10

READS

756

5 authors, including:



**Richard Witte**

Office National de la Chasse et de la Faune Sauvage

26 PUBLICATIONS 110 CITATIONS

[SEE PROFILE](#)



**Gerard van Buurt**

32 PUBLICATIONS 1,339 CITATIONS

[SEE PROFILE](#)



**Adolphe Oscar Debrot**

Wageningen University & Research

253 PUBLICATIONS 3,270 CITATIONS

[SEE PROFILE](#)



**Luis Bermúdez-Villapol**

Cetacean Research center. Venezuela

18 PUBLICATIONS 178 CITATIONS

[SEE PROFILE](#)

# First record of Fraser's dolphin *Lagenodelphis hosei* for the Dutch Caribbean

RICHARD H. WITTE<sup>1</sup>, GERARD VAN BUURT<sup>2</sup>, ADOLPHE O. DEBROT<sup>1</sup>, LUIS A. BERMUDEZ-VILLAPOL<sup>3</sup>  
AND FERNANDO SIMAL<sup>4</sup>

<sup>1</sup>IMARES, Wageningen University Research Centre, PO Box 57, 1780 AB Den Helder, The Netherlands, <sup>2</sup>Island Territory of Curaçao, Department of Agricultural, Animal Husbandry and Fisheries, Curaçao, Santa Rosaweg 122, Curaçao, <sup>3</sup>Centro de Investigación de Cetáceos CIC (Cetacean Research Center), E/S Los Robles, Isla de Margarita, 6313, Venezuela, <sup>4</sup>Natural and Historic Resources Unit, STINAPA Bonaire, PO Box 368, Bonaire

*A dead dolphin found on Bonaire in August 2011 is identified as adult Fraser's dolphin Lagenodelphis hosei, a new species for the Dutch Caribbean. A first closer examination showed a collapsed lung, stomach parasite infection and abundant mouth ulceration as indications of its health status. The animal was relatively fresh and did not die very long before it was found. Like more often with stranded deep diving cetacean species within the area, remnants of crustacean were found in its beak indicating recent foraging.*

**Keywords:** Fraser's dolphin, *Lagenodelphis hosei*, stranding, distribution, Bonaire, Leeward Antilles, Crustacean

Submitted 12 January 2012; accepted 14 February 2012

## INTRODUCTION

On 5 August, 2011, a dead dolphin was found on the bottom of Lac Bay, Bonaire in approximately 3 m of water but very close to the deeper channel, about 300 m from shore. Based on diagnostic photographs, the specimen could be identified as an adult Fraser's dolphin, *Lagenodelphis hosei* Fraser 1956, which means a first record for the Dutch Caribbean. As the species is exceedingly rare in the southern Caribbean, this paper documents the find and places the record within the regional context.

The Fraser's dolphin *Lagenodelphis hosei* Fraser 1956, was identified by its length (about 2.56 cm), a centrally placed slightly falcate but almost triangular dorsal fin of 19 cm in height (snout to anterior insertion of dorsal fin 109 cm) (e.g. Figure 2), short stubby beak (snout to blowhole 33 cm), dark at the upper side and pinkish lower jaw, short flippers, pointed to the tips (28 cm in length), dark grey back and pinkish abdomen (Figure 4) and broad flipper stripe that starts at the lower jaw (e.g. Figure 1). The fluke width was measured as 55 cm and the maximum girth as 117 cm. The true coloration was already faded, and the body's stiffness indicated that it had died at least several hours before it was found. The lateral band between snout and anus is almost indistinct.

Closer examination of the exterior by members of STINAPA Bonaire showed there was damage on the tip of the snout most likely caused by the dragging. Besides a clear small cookie-cutter shark-bite (likely *Isistius brasiliensis*) (e.g. Figure 3), scars caused by squids and several superficial scars and bruises, no other wounds were visible. The mouth, throat and blowhole were internally checked for objects with

negative results. Between the teeth were remnants of crustacean (Figure 5) which still need to be identified but that indicate recent foraging. Further, abundant mouth ulceration, stomatitis type, was found leading to tissue necrosis, with deformations at the edge of the palate and gums. The first stomach compartment contained stomach parasites (possibly of the genus *Anisakis*) and traces of algae. The mucosa in the second compartment was very red and haematized. The lungs were collapsed, showing abnormal consistency, with obvious wasting (invasive bleedings) mechanically blocking the respiratory function. After measuring and tissue sampling of liver, pancreas, lungs, blubber and collection of stomach parasites which were still living, the cadaver was discarded north-west of the island, close to Nukove.

## DISCUSSION

The Fraser's dolphin *Lagenodelphis hosei* was first described in 1956 based on a single skull found prior to 1895 by Charles Hose on a beach in Sarawak, Borneo (Fraser, 1956). However, it was not documented in the wild until 1971 (Perrin *et al.*, 1973). The extent of the geographical range is poorly known but based on strandings and sightings by now the species is believed to have a cosmopolitan distribution from approximately 30° south to 30° north, and to live in deep (usually over 1000 m deep), tropical waters. Fraser's dolphins have been observed in the northern Gulf of Mexico during all seasons where they occur in oceanic waters (>200 m) (Perrin *et al.*, 1994; Mullin & Hoggard, 2000). It appears they can occur in areas near coastlines where the continental shelf is narrow (Jefferson *et al.*, 2008).

Fraser's dolphins are, when seen, usually found in large, tight groups averaging between 10 and 100 individuals, but have been occasionally seen in larger groups of up to 1000

**Corresponding author:**

R.H. Witte

Email: richard.witte@wur.nl



Fig. 1. Lateral view of the Fraser's dolphin found on August 5 2011, Sorobon, Bonaire. Shape of snout, pectoral fin, dorsal fin, position of dorsal fin and coloration are clearly visible. © Gerard van Buurt.




Fig. 2. Dorsal fin close-up. © Gerard van Buurt.



Fig. 3. Mark of cookie-cutter shark bite. © Gerard van Buurt.



Fig. 4. Ventral view of the Fraser's dolphin found on Bonaire. The urogenital/anal opening is clearly visible as well as abdominal superficial scars and bruises. © Gerard van Buurt.




Fig. 5. Remnants of possibly *Gnatophausia* sp. found between the teeth.

or more animals. They are relatively often seen in mixed schools associated with other deep diving cetacean species such as false killer whales, melon-headed whales, Risso's dolphins, and short-finned pilot whales, all species observed in the vicinity of the Dutch Leeward Islands.

Fraser's dolphins are capable of diving up to almost 2000 feet (600 m) to feed on deep-sea species of mesopelagic fish (e.g. myctophids), crustaceans, and cephalopods (e.g. squid and octopus). *Gnatophausia* sp. shrimp for instance are common in the stomachs of Caribbean pelagic cetaceans, especially beaked whales (ziphiids) (Debrot & Barros, 1992; Rosario-Delestre & Mignucci-Giannoni, 1998; Rosario-Delestre *et al.*, 1999) and pygmy sperm whale (*Kogia breviceps*) (Secchi *et al.*, 1994). The remnants that were found in its beak (possibly *Gnatophausia* sp.) suggest that deep water mysid may be also an important prey for Fraser's dolphins.

In the past decade several studies have become available which provide additional information on the marine mammal fauna of the leeward Dutch waters (e.g. Debrot & Barros, 1994; Debrot, 1998; Debrot *et al.*, 1998, 2006, 2011; Maldani, 2008; Luksenburg, 2011). However, most records have remained uncompiled and unassessed and therefore Debrot *et al.* (in press) recently provided an assessment of the marine mammals of the windward Dutch islands and Witte *et al.* (unpublished data) of the leeward Dutch island. Based on that it became clear that this was the first Fraser's dolphin *Lagenodelphis hosei* (Fraser, 1956), found in the waters of the former Dutch Antilles.

The IUCN Red List of Threatened Species considers Fraser's dolphins as 'Data Deficient' due to insufficient information on population status and trends. The reported number of sightings of Fraser's dolphins in the Atlantic Ocean are in the low tens (Hersh & Odell, 1986; Ward *et al.*, 2001) and just over a dozen for the Caribbean Sea. As the species is very rare, all records should be reported and documented as fully as possible. The Caribbean records include three specimens harpooned and taken by a cetacean fishery at St Vincent between 1972 and 1973 (Caldwell *et al.*, 1976), two sightings off Dominica on 26 and 28 October 1991 (Watkins *et al.*, 1994) and two strandings records in Puerto Rico: one subadult male on the south-west coast in 1994, and a female calf in 1997 in Ponce, on the south coast (Mignucci-Giannoni *et al.*, 1999). Carlson *et al.* (1995) reported sightings of Fraser's dolphins off the leeward coast of Dominica during January, February, and March of 1995, and IFAW (1996) reported sightings offshore of Carriacou in 1996. Two dolphins live stranded on El Palito Beach, State of Carabobo, Venezuela, on 6 June 1999. This was the first record of this species for Venezuela and the southern Caribbean (Bolaños & Villarroel-Marín, 2003). Boisseau *et al.*

(2000) reported two new encounters off Dominica: one pair and a group of approximately 50 individuals. Based on two sightings (130 individuals) during surveys off La Martinique in March and April 2003 they are now considered common in its territorial waters (Jeremie, 2005).

Based on this stranding and other reported sightings it can be concluded that within the Caribbean Sea, Fraser's dolphins are likely to be found in deeper waters, both offshore and near shore. On islands with near shore deep waters occasionally they can be found stranded. Within the Dutch Caribbean waters this species is most likely to be found in offshore deeper waters. The presence of krill (*Euphausia* sp.) may play a significant role in the distribution and species richness of cetaceans in the Caribbean Sea and should be further investigated.

Positive determination at sea is not easy, especially for inexperienced observers, and confusion with other dolphin species is not unlikely. Therefore it is recommended to always photograph or even film any sightings. Especially in the case of dolphins found within a group of 'black-fish' such as short-finned pilot whales or melon-headed whales, one should be on the lookout for Fraser's dolphin.

In order to establish which species are present, provide evidence of seasonal residency patterns and to stimulate people to report their sightings, these updated reviews were made and all known sightings were compiled into the Dutch Caribbean Marine Mammal Database (DC-MMDB) (Witte *et al.*, unpublished data). To help stimulate the general public to report their encounters with marine mammals, the records compiled up to now will be added to the public data website ([www.observado.org](http://www.observado.org)).

## ACKNOWLEDGEMENTS

This paper is dedicated to the memory of Dr P.J.H. van Bree curator of mammals in the Zoological Museum of the University of Amsterdam who passed away on 24 February 2011. Dr van Bree conducted a lot of systematic work on marine mammals (in particular cetaceans and pinnipedia) and he was the first to make a preliminary list of the cetaceans of the southern Caribbean (van Bree, 1975). We thank Elsmarie Beukeboom (STINAPA Bonaire), Jaime Bolanos (SEA VIDA), Paul Hoetjes (EL&I and SCCN) and Ramon de Leon (STINAPA Bonaire) for their information and cooperation and of course remain especially indebted to the finders. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

## REFERENCES

- Boisseau O., Carlson C.A., and Seipt I. (2000) *A report on cetacean research conducted by the International Fund for Animal Welfare (IFAW) off Guadeloupe, Dominica, Martinique, Grenada and Tobago from 12 January to 30 March 2000*. Unpublished Report to the International Fund for Animal Welfare.
- Bolaños J. and Villarroel-Marín A. (2003) Three new records of cetacean species for Venezuelan waters. *Caribbean Journal of Science* 39, 230–232.
- van Bree P.J. (1975) Preliminary list of the cetaceans of the southern Caribbean. *Studies on the Fauna of Curaçao and other Caribbean Islands* 48, 79–87.
- Debrot A.O. (1998) New cetacean records for Curaçao, Netherlands Antilles. *Caribbean Journal of Science* 34, 168–170.
- Debrot A.O. and Barros N.B. (1992) Notes on a Gervais' beaked whale, *Mesoplodon europaeus*, and a dwarf sperm whale, *Kogia simus*, stranded in Curaçao, Netherlands Antilles. *Marine Mammal Science* 8, 172–178.
- Debrot A.O. and Barros N.B. (1994) Additional cetacean records for the leeward Dutch Antilles. *Marine Mammal Science* 10, 359–368.
- Debrot A.O., de Meyer J.A. and Dezentjé P.J.E. (1998) An overview of the cetacean fauna of the leeward Dutch Antilles. *Caribbean Journal of Science* 34, 204–210.
- Debrot A.O., van Buurt G., Caballero A. and Antczak A.A. (2006) A historical review of records of the West Indian manatee and the American crocodile in the Dutch Antilles. *Caribbean Journal of Science* 42, 272–280.
- Debrot A.O., Witte R.H. and Scheidat M. (2011) *The marine mammals of the Dutch Caribbean: a comparison between EEZ sectors, contrasts and concerns*. Paper presented to IWC Scientific Committee SC/63/E9, 8 pp.
- Debrot A.O., Esteban N., Bervoets T., Hoetjes P.C. and Scheidat M. (in press) Marine mammals of the north-eastern Caribbean Windward Dutch Islands: Saba, St Eustatius, St Maarten, and the Saba Bank. *Caribbean Journal of Science*.
- Caldwell D.K., Caldwell M.C. and Walker R.V. (1976) First records for Fraser's dolphin (*Lagenodelphis hosei*) in the Atlantic and melon-headed whales (*Peponocephala electra*) in the W. Atlantic. *Cetology* 25, 1–4.
- Carlson C.A., Seipt I., Brown R., Lewis E. and Moscrop A. (1995) *Report on a project by the International Fund for Animal Welfare (IFAW) to enhance public awareness and promote the appropriate development of whale watching in Dominica*. International Whaling Commission, an Information Paper, Working Group on Whale Watching, 15 pp.
- Fraser F.C. (1956) A new Sarawak dolphin. *Sarawak Museum Journal* 7, 478–503.
- Hersh S.L. and Odell D.K. (1986) Mass strandings of Fraser's dolphin, *Lagenodelphis hosei*, in the western North Atlantic. *Marine Mammal Science* 2, 73–76.
- IFAW (International Fund for Animal Welfare) (1996) *Cetacean field research conducted from song of the whale off Dominica and Grenada: spring 1996*. Unpublished Report to the International Fund for Animal Welfare.
- Jefferson T.A., Webber M.A. and Pitman R.L. (2008) *Marine mammals of the world: a comprehensive guide to their identification*. Amsterdam: Elsevier.
- Jeremie S. (2005) *Synthesis of cetaceans off the island of Martinique, French West Indies*. Regional Workshop of Experts on the Development of the Marine Mammal Action Plan for the Wider Caribbean Region, Bridgetown, Barbados.
- Luksenburg J.A. (2011) Three new records of cetacean species for Aruba, Leeward, Antilles, southern Caribbean. *Marine Biodiversity Records* 4, 1–4. DOI: 10.1017/S1755267210001193.
- Maldani D. (2008) *Preliminary observations of bottlenose dolphins around Bonaire, Netherland Antilles*. Kralendijk, Bonaire, Netherland Antilles: Council for International Exchange, Research Station Bonaire.
- Mignucci-Giannoni A.A., Montoya-Ospina R.A., Pérez-Zayas J.J., Rodríguez-López M.A. and Williams E.H. Jr (1999) New records of Fraser's dolphin (*Lagenodelphis hosei*) for the Caribbean. *Aquatic Mammals* 25, 15–19.
- Mullin K.D. and Hoggard W. (2000) Visual surveys of cetaceans and sea turtles from aircraft and ships. In Davis R.W., Evans W.E. and Würsig



B. (eds) *Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: distribution, abundance and habitat associations. Volume II: Technical report. OCS Study MMS 96-0027*. New Orleans, LA: Minerals Management Service, Gulf of Mexico OCS Region, pp. 111–172.

**Perrin W.F., Best P.B., Dawbin W.H., Balcomb K.C., Gambell R. and Ross G.J.B.** (1973) Rediscovery of Fraser's dolphin (*Lagenodelphis hosei*). *Nature* 241, 345–350.

**Perrin W.F., Leatherwood S. and Collet A.** (1994) Fraser's dolphin, *Lagenodelphis hosei* Fraser, 1956. In Ridgway S.H. and Harrison R. (eds) *Handbook of marine mammals. Volume 5. The first book of dolphins*. London: Academic Press, pp. 225–240.

**Rosario-Delestre R.J. and Mignucci-Giannoni A.A.** (1998) Epizootic and herd strandings of goosebeak whales in the Caribbean. In *Abstracts of the World Marine Mammal Science Conference (Society for Marine Mammalogy)*, January 1998, Monaco, p. 116.

**Rosario-Delestre R.J., Rodríguez-López M.A., Mignucci-Giannoni A.A. and Mead J.G.** (1999) New records of beaked whales

(*Mesoplodon* spp.) for the Caribbean. *Caribbean Journal of Science* 35, 144–148.

**Secchi E.R., Campolim M.B. and Möller L.M.** (1994) Notas sobre o enalhe de dois cachalotes pigmeus *Kogia breviceps* na costa sul do Rio Grande do Sul, Brasil. In *Actas, IV Reunión de Trabajo de Especialistas en Mamíferos Acuáticos de América del Sur*, 12–15 November 1990, Valdivia, Chile, pp. 244–262.

and

**Watkins W.A., Daher M.A., Frstrup K. and Notarbartolo-di-Sciara G.** (1994) Fishing and acoustic behaviour of Fraser's dolphin (*Lagenodelphis hosei*) near Dominica, southeast Caribbean. *Caribbean Journal of Science* 30, 76–82.

#### Correspondence should be addressed to:

R. Witte  
IMARES, Wageningen University Research Centre  
PO Box 57, 1780 AB Den Helder, The Netherlands  
email: richard.witte@wur.nl